



3 1761 05289038 1

Digitized by the Internet Archive
in 2008 with funding from
Microsoft Corporation




24

2290

7

ILLUSTRATIONS
OF
BRITISH FUNGI.



Bot
halloph
C

ILLUSTRATIONS
OF
BRITISH FUNGI
(HYMENOMYCETES),

To serve as an Atlas to the "Handbook of British Fungi."

BY

M. C. COOKE, M.A., LL.D., A.L.S.,

President of the Quekett Microscopical Club, and of the Hackney Natural History and Microscopical Society. Fellow of the Botanical Society of Edinburgh. Corresponding Member of the Academy of Natural Sciences, Philadelphia; of the Academy of Science of New York; of the Portland Society of Natural History, Maine; and of the Società Crittogamologica Italiana. Honorary Member of the Cryptogamic Society of Scotland; the Woolhope Field Club; the Hertfordshire Natural History Society; the Norfolk and Norwich Naturalists' Society; the Essex Field Club; the Birmingham Natural History Society; and other Societies

VOL. VII.

137427
611/16

London:

WILLIAMS AND NORGATE, HENRIETTA STREET, W.C.

1888-1890.

QK

607

C67

V.7

SYSTEMATIC INDEX.

GEN. 9. **LACTARIUS**, *Fr. Epic.* 333.

Hymenophore continuous with the stem. Gills unequal, between membranaceous and waxy, rather rigid, *milky*, edge acute. Spores globose, white, rarely turning yellowish. Fleshy, terrestrial, putrescent fungi, with a depressed pileus, and adnate-decurrent, sometimes branched gills.

Tribe 1. Piperites. Stem central. Gills unchangeable, naked, not discoloured or pruinose. Milk at first white, commonly acid.

* **TRICHOLOMOIDEI.** Pileus moist, viscid, margin at first involute and tomentose.

922.	serobiculatus, <i>Scop.</i>	.	.	971
923.	torminosus, <i>Schæff.</i>	.	.	972
924.	cilicioides, <i>Fries</i>	.	.	973
—	intermedius, <i>Kromb.</i>	.	.	—
925.	turpis, <i>Weinm.</i>	.	.	987
—	var. plumbeus, <i>Fries.</i>	.	.	—
926.	controversus, <i>Pers.</i>	.	.	1003
927.	pubescens, <i>Fries</i>	.	.	974
928.	aspideus, <i>Fries</i>	.	.	1083

** **LIMACINI.** Pileus moist, viscid, with a pellicle; margin naked.

929.	insulsus, <i>Fries</i>	.	.	975
—	zonarius, <i>Bull</i>	.	.	—
930.	utilis, <i>Weinm.</i>	.	.	1084
931.	blennius, <i>Fries</i>	.	.	988
932.	hysginus, <i>Fries</i>	.	.	989
933.	trivialis, <i>Fries</i>	.	.	976
934.	circellatus, <i>Batt.</i>	.	.	990
935.	uvidus, <i>Fries</i>	.	.	991

* * **PIPERATI.** Pileus without pellicle, hence absolutely dry, often unpolished.

936.	flexuosus, <i>Fries</i>	.	.	992
937.	pyrogalus, <i>Bull</i>	.	.	993
938.	squalidus, <i>Kromb.</i>	.	.	1004a
939.	capsicum, <i>Schulz.</i>	.	.	977
940.	chrysorrheus, <i>Fries</i>	.	.	984
941.	acris, <i>Bolt.</i>	.	.	1005

942.	umbrinus, <i>Pers.</i>	.	.	1006
943.	pargamenus, <i>Swartz.</i>	.	.	978
944.	piperatus, <i>Scop.</i>	.	.	979
945.	vellerens, <i>Fries</i>	.	.	980
946.	exsuccus, <i>Otto.</i>	.	.	981
938.	scoticus, <i>B. & Br.</i>	.	.	1004b

Tribe 2. Dapetes. Stem central. Gills naked, milk at first brightly coloured.

947.	deliciosus, <i>Linn.</i>	.	.	982
------	--------------------------	---	---	-----

Tribe 3. Russularia. Stem central. Gills pallid, then discoloured, then darker, changing with the light, at length pruinose with white. Milk at first white, either mild, or mild then becoming acrid.

* *Pileus* at first viscid.

948.	pallidus, <i>Pers.</i>	.	.	1007
949.	quietus, <i>Fries</i>	.	.	983
950.	aurantiacus, <i>Fl. Dan.</i>	.	.	1099
—	theiogalus, <i>Bull</i>	.	.	—
951.	cremor, <i>Fries</i>	}	.	1008
	var. pauper, <i>Karst.</i>			
952.	vietus, <i>Fries</i>	.	.	1009
953.	cyathula, <i>Fries</i>	.	.	1085

** *Pileus* unpolished, squamulose, villous, or pruinose.

954.	rufus, <i>Scop.</i>	.	.	985
955.	helvus, <i>Fries</i>	.	.	994
956.	tomentosus, <i>Otto.</i>	.	.	1010
—	mammosus, <i>Fries</i>	.	.	—
957.	var. monstrosus, <i>Fr.</i>	.	.	995
958.	glyciosmus, <i>Fries</i>	.	.	1011
959.	fuliginosus, <i>Fries</i>	.	.	996
960.	picinus, <i>Fries</i>	.	.	997
961.	{ lilacinus, <i>Lasch.</i>	.	.	998a
	{ spinosulus, <i>Quel.</i> var.	.	.	998b

*** *Pileus* polished, smooth.

962.	volemus, <i>Fries</i>	.	.	999
963.	ichoratus, <i>Batsch.</i>	.	.	1000
964.	serifluus, <i>D. Cand.</i>	.	.	1012
965.	mitissimus, <i>Fries</i>	.	.	1001
966.	subdulcis, <i>Bull</i>	.	.	1002
—	var. Terrei, <i>B & Br.</i>	.	.	—
967.	{ camphoratus, <i>Fries</i>	.	.	1013a
	{ cimicarius, <i>Batsch.</i>	.	.	1013b
968a.	subumbonatus, <i>Lind.</i>	.	.	986a
969a.	obnubilis, <i>Lasch.</i>	.	.	1014a
968b.	minimus, <i>Smith</i>	.	.	986b

Tribe 4. Pleuropus. Stem excentric or lateral.

969*b*. obliquus, *Fries*. . . 1014*b*

GEN. 10. **RUSSULA**, *Pers.*

Veil none. Hymenophore descending unchanged into the vesiculose trama. Gills rigid, fragile, not milky, edge acute. Spores rounded, often echinulate, white, or becoming yellow. Terrestrial, fleshy, putrescent fungi, stem polished, pileus at first or at length depressed.

- I. **COMPACTÆ.** Pileus everywhere fleshy, hence the margin at first turned in, always without striæ, or a distinct viscid pellicle (hence the colour not variable, only changing with age or the action of the air), flesh firm, compact. Stem solid, fleshy. Gills unequal.

970.	nigricans, <i>Bull.</i>	.	1015
971.	var. albo-nigra, <i>Krom.</i>	.	1016
972.	adusta, <i>Pers.</i>	.	1051
973.	densifolia, <i>Secr.</i>	.	1017
974.	semicrema, <i>Fries</i>	.	1067
975.	delica, <i>Fries</i>	.	1068
—	elephantina, <i>Fries</i>	.	?
976.	mustelina, <i>Fries</i>	.	1018

- II. **FURCATÆ.** Pileus compact, firm, covered with a thin, closely adnate pellicle, which at length disappears, margin abruptly thin, at first turned in, then spreading, acute, even. Stem at first compact, at length soft and spongy within. Gills rather forked, mixed with a few shorter ones, commonly attenuated towards each end, thin and normally narrow.

977.	olivascens, <i>Fries</i>	.	1035
978.	furcata, <i>Pers.</i>	.	1036
979.	var. pictipes, <i>Cke.</i>	.	1086
980.	var. ochroviridis, <i>Cke.</i>	.	1100
981.	sanguinea, <i>Bull</i>	.	1019
982.	rosacea, <i>Fries</i>	.	1020
983.	maculata, <i>Quel.</i>	.	1069
984.	sardonina, <i>Fries</i>	.	1037
985.	depallens, <i>Fries</i>	.	1021
986.	purpurea, <i>Gillet</i>	.	1022
987.	cærulea, <i>Pers.</i>	.	1052
988.	drimæia, <i>Cooke</i>	.	1023

- III. **RIGIDÆ.** Pileus destitute of a viscid pellicle, absolutely dry, rigid, the cuticle commonly breaking up into flocci or granules. Flesh

thick, compact, firm, vanishing near the margin, which is straight (never involute), spreading, and without striæ. Stem solid, at first hard, then softer and spongy. Of the gills, a few reach half-way to the stem; the others are divided, rigid, dilated in front, and extend into a very broad, rounded end, whence the margin of the pileus becomes obtuse and not inflexed.

989.	<i>lactea</i> , <i>Pers.</i>	. . .	1070
990.	var. <i>incarnata</i>	. . .	1071
991.	<i>virescens</i> , <i>Schæff.</i>	. . .	1039
992, 993.	<i>cutefracta</i> , <i>Cooke</i>	. . .	1024, 1040
994, 995.	<i>lepida</i> , <i>Fries</i>	. . .	1072, 1073
996.	<i>rubra</i> , <i>D. Cand.</i>	. . .	1025
997.	var. <i>atropurpurea</i> , <i>Kromb.</i>	. . .	1087
998.	<i>Linnæi</i> , <i>Fries</i>	. . .	1026
999, 1000.	<i>xerampelina</i> , <i>Schæff.</i>	. . .	1053, 1074
1001.	<i>olivacea</i> , <i>Schæff.</i>	. . .	1041

IV. HETEROPHYLLÆ. Pileus fleshy, firm, covered with a thin adnate pellicle, margin thin, at first turned in, then expanded and striate. Gills many shorter mixed with the longer ones, others forked. Stem solid, stout, spongy within.

1002.	<i>vesca</i> , <i>Fries</i>	. . .	1075
1003b.	<i>Du Portii</i> , <i>Phil.</i>	. . .	1042b
1004.	<i>lilacea</i> , <i>Quelet</i>	. . .	1054
1005.	<i>azurea</i> , <i>Bres.</i>	. . .	1088
1003a.	<i>serotina</i> , <i>Quelet</i>	. . .	1042a
1006, 1007, 1008.	<i>cyanoxantha</i> , <i>Schæff.</i>	1043, 1076, 1077	
1009, 1010.	<i>heterophylla</i> , <i>Fries</i>	. . .	1044, 1045
1011.	<i>galochroa</i> , <i>Bull</i>	. . .	1089
1012.	<i>consobrina</i> , <i>Fries</i>	. . .	1055
1013.	var. <i>intermedia</i> , <i>Cke.</i>	. . .	1056
1014.	var. <i>sororia</i> , <i>Larb.</i>	. . .	1057
1015.	<i>foetens</i> , <i>Pers.</i>	. . .	1046
1016.	<i>subfoetens</i> , <i>Smith</i>	. . .	1047
1017.	<i>fellea</i> , <i>Fries</i>	. . .	1058
1018.	<i>elegans</i> , <i>Bres.</i>	. . .	1027
1019.	<i>Quelettii</i> , <i>Fries</i>	. . .	1028
1020.	<i>expallens</i> , <i>Gillet</i>	. . .	1029

V. FRAGILES. Pileus more or less fleshy, rigid, but fragile, covered with a pellicle, always continuous, viscid after rain, and somewhat separable, margin membranaceous, at first closed in, but not involute, when adult commonly sulcate and

tuberculose. Flesh usually floccose, lax and friable. Stem spongy, at length soft and hollow. Gills almost all equal, simple, broadest in front, free in the pileus when closed.

* Gills and spores white.

1021.	<i>emetica, Fries</i>	.	.	1030
1022.	var. <i>clusii, Vitt.</i>	.	.	1031
1023.	var. <i>fallax, Schæff.</i>	.	.	1059
1024.	<i>pectinata, Bull.</i>	.	.	1101
1025.	<i>ochroleuca, Pers.</i>	.	.	1049
1026.	<i>granulosa, Cooke</i>	.	.	1038
1027.	<i>æruginea, Fries</i>	.	.	1090
1028.	<i>fragilis, Pers.</i>	.	.	1091
1029.	{ var. <i>violacea, Fr.</i>	.	.	1060a
	var. <i>nivea, Fr.</i>	.	.	1060b
1030.	<i>figibilis, Britz.</i>	.	.	1048
1031.	<i>citrina, Gillet</i>	.	.	1078
1032.	<i>punctata, Gillet</i>	.	.	1032

** Gills and spores white, then yellow or bright lemon.

1033, 1034.	<i>veternosa, Fries</i>	.	.	1033, 1092
1035.	var. <i>roseipes, Bres.</i>	.	.	1081
1036, 1037.	<i>integra, Linn.</i>	.	.	1034, 1093
1038.	var. <i>alba, Cke.</i>	.	.	1094
1039.	<i>decolorans, Fries</i>	.	.	1079
1040.	<i>Barlx, Quelet.</i>	.	.	1061
1041.	<i>aurata, With.</i>	.	.	1080
1042, 1043.	<i>nitida, Pers.</i>	.	.	1062, 1063
1044.	{ var. <i>cuprea, Krom.</i>	.	.	1095b
	<i>pulchralis, Britz.</i>	.	.	1095a
1045.	<i>armeniaca, Cooke</i>	.	.	1064
1046, 1047.	<i>puellaris, Fries</i>	.	.	1065, 1066

* * * Gills and spores ochre.

1048, 1049.	<i>alutacea, Fries</i>	.	.	1096, 1097
1050.	<i>ochracea, A. & S.</i>	.	.	1050
1051.	<i>lutea, Huds.</i>	.	.	1082
1052.	<i>nauseosa, Pers.</i>	.	.	1102a, 1147
1053.	<i>vitellina, Pers.</i>	.	.	1102b
1054.	<i>chameleontina, Fries</i>	.	.	1098

GEN. 11. **CANTHARELLUS**, *Adans.*

Hymenophore continuous with the stem, descending unchanged into the trama. Gills thick, between fleshy and waxy, fold-like, rather branched, edge obtuse. Spores white. Fleshy membranaceous putrescent fungi, veil none.

1. **MESOPUS.** Pileus entire. Stem central.

* Pileus and solid stem fleshy.

1055.	<i>cibarius</i> , <i>Fries</i> .	. 1103
1056.	{ var. <i>rufipes</i> .	. 1131a
	{ <i>Friesii</i> , <i>Quelet</i> . .	. 1131b
1057.	<i>aurantiacus</i> , <i>Wulf.</i> .	. 1104
1058a.	<i>Brownii</i> , <i>B. & Br.</i> .	. 1106a
1059.	<i>carbonarius</i> , <i>A. & S.</i> .	} . 1105
	var. <i>radicosus</i> , <i>B. & Br.</i> }	
1058b.	<i>umbonatus</i> , <i>Fries</i> .	. 1106b
1060a.	<i>albidus</i> , <i>Fries</i> .	. 1107a

** Pileus submembranaceous, stem hollow, polished.

1060b.	<i>Houghtoni</i> , <i>Phil.</i> .	. 1107b
1061.	<i>tubæformis</i> , <i>Fries</i> .	. 1108
—	var. <i>lutescens</i> , <i>Bull.</i> .	—
1062.	<i>infundibuliformis</i> , <i>Fries</i> .	. 1109
1063a.	<i>cinereus</i> , <i>Pers.</i> .	. 1110a
1064a.	<i>leucophæus</i> , <i>Nouel.</i> .	. 1111a
1063b.	<i>cupulatus</i> , <i>Fries</i> .	. 1110b
1098a.	<i>devexus</i> , <i>Fries</i> .	. 1150a
1064b.	<i>Stevensoni</i> , <i>B. & Br.</i> .	. 1111b

II. PLEUROPUS. Dimidiate, stem lateral.

1065.	{ <i>muscigenus</i> , <i>Bull.</i> .	. 1115a
	{ <i>glaucus</i> , <i>Fries</i> .	. 1115b

III. RESUPINATI. Pileus entire, at first cup-shaped, fixed at the vertex, then subreflexed.

* *Bryophili.*

1066.	{ <i>retirugus</i> , <i>Bull.</i> .	. 1112a
	{ <i>lobatus</i> , <i>Fries</i> .	. 1112b

GEN. 12. **NYCTALIS**, *Fr. Syst. Orb.* 203.

Hymenophore continuous with the stem. Gills fleshy, thick, juicy, edge obtuse, not decurrent on the stem nor fold-like. Veil floccosely pruinose. Fleshy fungi, not reviving, of uncertain occurrence.

I. SPELEÆ. Gills crowded, somewhat growing together.

1067a.	<i>caliginosa</i> , <i>Smith</i> .	. 1132a
--------	------------------------------------	---------

II. PARASITÆ. Gills distinct, distant, growing on putrid fungi.

1067b.	<i>asterophora</i> , <i>Fries</i> .	. 1132b
1068.	<i>parasitica</i> , <i>Bull.</i> .	. 1113

GEN. 13. **MARASMIUS**, *Fr. Gen. Hym.* 372.

Tough, dry, shrivelling (not putrescent), fungi reviving when moistened. Hymenophore continuous with the stem,

but heterogeneous, descending into the trama; veil none. Stem cartilaginous or horny. Gills tough, rather distant, edge acute and entire.

- I. COLLYBIA. Pileus fleshy, tough, at length rather leathery, sulcate or wrinkled, margin at first turned in. Stem somewhat cartilaginous. Mycelium floccose, or in a few not manifest.

- A. SCORTEI. Stem solid, or stuffed, with a pith, then hollow, fibrous within, externally with a villous down, easily wiped away, which covers the cartilaginous cuticle. Gills receding, free.

- * Stem woolly or strigose at the base.

1069.	urens, <i>Bull.</i>	.	.	1116
1070.	peronatus, <i>Bolt.</i>	.	.	1117
1071.	porreus, <i>Pers.</i>	.	.	1133

- ** Stem naked at the base, often interwoven with twisted fibres.

1072.	oreades, <i>Bolt.</i>	.	.	1118
1073.	{ plancus, <i>Fries</i>	.	.	1119a
	{ scorteus, <i>Fries</i>	.	.	1119b

- B. TERGINI. Stem rooting, definitely tubular, not fibrous, manifestly cartilaginous. Gills receding, free. Pileus thinner than in the preceding, hygrophanous, smooth, or striate at the margin.

- * Stem woolly below, smooth above.

1074.	prasiosmus, <i>Fries</i>	.	.	1120
1075.	{ varicosus, <i>Fries</i>	.	.	1121a
	{ fuscopurpureus, <i>Pers.</i>	.	.	1121b
1076a.	terginus, <i>Fries</i>	.	.	1122a

- ** Stem everywhere pruinat, velvety.

1077.	{ Wynnei, <i>B. & Br.</i>	.	.	1123a
	{ erythropus, <i>Fries</i>	.	.	1123b
1076b.	archyropus, <i>Pers.</i>	.	.	1122b
1078.	{ torquescens, <i>Quel.</i>	.	.	1124a
	{ impudicus, <i>Fries</i>	.	.	1124b

- C. CALOPODES. Stem short, not rooting, grafted, often with a floccose tubercle at the base. Pileus convex, involute, then plane and depressed, in which state the gills, typically adnate, are somewhat decurrent.

- * Stem very smooth and shining above, simple at the base.

1079.	{ scorodonius, <i>Fries</i>	.	.	1125a
	{ calopus, <i>Pers.</i>	.	.	1125b

1080.	{	vaillantii, <i>Fries</i>	.	.	1126a
		angulatus, <i>Pers.</i>	.	.	1126b
		languidus, <i>Fries</i>	.	.	1126c

** Stem velvety or pruinose, base rather tuberculose.

1081a.	fœtidus, <i>Sow.</i>	.	.	1134a
1082a.	amadelpus, <i>Fries</i>	.	.	1127a
1082.	{	ramealis, <i>Bull.</i>	.	1127b
		candidus, <i>Bolt.</i>	.	1127c

Tribe 2. Mycena. Stem horny, fistulose, sometimes medullate, tough, dry, mycelium rhizomorphoid, corticate not floccose. Pileus rather membranaceous, campanulate, then expanded, margin at first straight, pressed to the stem.

A. CHORDALES. Stem rigid, rooting, or adnate by a dilated base. Pileus campanulate or convex.

1083.	{	alliaceus, <i>Jacq.</i>	.	.	1128a
		cohærens, <i>Pers.</i>	.	.	1128b
1081b.		cauticinalis, <i>With.</i>	.	.	1134b

B. ROTULÆ. Stem filiform, flaccid, grafted at the base. Pileus soon rather plane or umbilicate.

* Stem quite smooth, shining.

1084.	{	rotula, <i>Scop.</i>	.	.	1129a
		graminum, <i>Lib.</i>	.	.	1129b
		androsaceus, <i>Linn</i>	.	.	1129c
1085.	{	splachnoides, <i>Fries</i>	.	.	1130a
		Curreyi, <i>B. & Br.</i>	.	.	1130b

** Stem velvety or pilose.

1085c.		perforans, <i>Fries</i>	.	.	1130c
1086.	{	institius, <i>Fries</i>	.	.	1135a
		Hudsoni, <i>Pers.</i>	.	.	1135b
1087.	{	epichlœ, <i>Fries</i>	.	.	1136a
		actinophorus, <i>B. & Br.</i>	.	.	1136b
		saccharinus, <i>Fries</i>	.	.	1136c
1088.	{	epiphyllus, <i>Fries</i>	.	.	1137a
		polyadelphus, <i>Lasch.</i>	.	.	1137b

Tribe 3. Apus. Pileus sessile, resupinate.

1088c.	spodoleucus, <i>Berk.</i>	.	.	} 1137c
	including <i>M Broomei</i> , <i>B.</i>	.	.	

GEN. 14. **LENTINUS**, *Fries.*

Pileus between fleshy and leathery, tough, hard when old, persistent. Hymenophore continuous with the stem, or base of the pileus when sessile. Gills concrete with the

hymenophore, thin, unequal, membranaceous, with the edge serrate or torn into teeth. Spores white, roundish, even. Growing on wood.

I. MESOPODES. Pileus nearly entire, stem distinct.

* *Lepidei*. Pileus squamose, more or less manifestly veiled.

1089.	tigrinus, <i>Bull.</i>	.	1138, 1139a
1090.	Dunalii, <i>D. Cand.</i>	.	1139b
1091, 1092.	lepidus, <i>Fries</i>	.	1140, 1141

** *Pulverulenti*. Pileus villose or pulverulent.

—	leontopodius, <i>Schultz</i>	.	—
—	pulverulentus, <i>Scop.</i>	.	—
—	resinaceus, <i>Trog.</i>	.	—
—	adhærens, <i>Alb. & Schw.</i>	.	—

*** *Cochleati*. Pileus smooth.

1093a.	cochleatus, <i>Fries</i>	.	1142a
--------	--------------------------	---	-------

II. PLEUROT. Dimidiate, sessile, or with a rather lateral stem.

1094.	scoticus, <i>B. & Br.</i>	.	1143
1095a.	fimbriatus, <i>Curr.</i>	.	1148a
1093b.	vulpinus, <i>Fries</i>	.	1142b
1095b.	flabelliformis, <i>Bolt.</i>	.	1148b

GEN. 15. **PANUS**, *Fries Epicr.* 396.

Whole fungus fleshy-leathery, tough, becoming dry, substance fibrous, radiating into the hymenium. Gills concrete with the hymenophore, unequal, at length leathery, edge entire. Spores even, white.

* Pileus irregular, stem excentric.

1096.	{ conchatus, <i>Fries</i>	.	1149a
	{ torulosus, <i>Fries</i>	.	1149b

** Stem definitely lateral.

1097.	{ stypticus, <i>Bull.</i>	.	1144a
	{ farinaceus, <i>Schum.</i>	.	1144b

*** Pileus resupinate, sessile, or extended behind.

1097c.	patellaris, <i>Fries</i>	.	1144c
—	Stevensoni, <i>B. & Br.</i>	.	—

GEN. 16. **XEROTUS**, *Fries Elen.* I. 48.

Hymenophore continuous with the stem, descending into the trama, which is similar to the leathery pileus. Gills leathery, broadly fold-like, dichotomous, edge entire, obtuse.

1089*b*. *degener*, *Fries* . . . 1150*b*

GEN. 17. **TROGIA**, *Fries Epic.* 402.

Gills fold-like, longitudinally channelled along the edge, in other respects like *Xerotus*.

1099*a* *crispa*, *Pers.* . . . 1114*a*

GEN. 18. **SCHIZOPHYLLUM**, *Fries Obs.* i. 103.

Pileus without flesh, dry. Gills leathery, branched in a fan-like manner, united above by the tomentose pellicle, bifid, edge longitudinally split. Spores roundish, white.

1099*b* *commune*, *Fries* . . . 1114*b*

GEN. 19. **LENZITES**, *Fries Epic.* 403.

Pileus corky or leathery, substance dry and floccose. Gills leathery, firm, either simple, unequal, or anastomosing and forming pores behind, trama floccose and similar to the pileus, edge rather acute.


* Growing on deciduous trees.

1100. { *betulina*, *Linn* . . . 1145*a*
 { *flaccida*, *Bull.* . . . 1145*b*


** Growing on conifers.

1101. { *sepiaria*, *Fries* . . . 1146*a*
 { *abietina*, *Bull.* . . . 1146*b*

milk white, then yellow.




LACTARIUS (PIPERITES) SCROBICULATUS. Scop.
on the ground, near Bristol.






LACTARIUS (PIPERITES) TORMINOSUS. Schæff.
on the ground in woods. Aymestry, Oct. 1871.





LACTARIUS (PIPERITES) CILICIOIDES. Fries.
on the ground. near Hereford.



milk white, acrid.




viscid.

8 μ

LACTARIUS (PIPERITES) TURPIS. *Weinm.*

in woods. Dinnorst, Oct. 1881.




milk white, acrid.



6-8/M


LACTARIUS (PIPERITES) CONTROVERSUS. Pers.
in pastures, &c. near Hereford.






LACTARIUS (PIPERITES) PUBESCENS. *Fries.*

in open grassy places. Ledbury, Oct. 1883.






LACTARIUS (PIPERITES) ASPIDEUS. *Fries.*
 in swampy places. Harewood, near Leeds. Sept. 1888.





LACTARIUS (PIPERATES) INSULSUS. *Fries.*

in woods. Mattishall, Oct. 1884.




milk white.

mild.



LACTARIUS (PIPERITES) UTILIS. *Weinm.*
on the ground. near Grove Park, Warwick, 1888.




milk white, acrid.

glutinous.

8 X 6 M

LACTARIUS (PIPERITES) BLENNIUS. *Fries.*
in beech woods. Epping, 1884.




milk white, acrid.

H.G.B.

8 x 10 M


LACTARIUS (PIPERITES) HYGGINUS. *Fries.*

amongst grass. Letton Court and Epping Forest, 1886.



milk white, acrid.


LACTARIUS (PIPERITES) TRIVIALIS. *Fries.*
in moist woods. near Hereford.



milk white, acrid.




LACTARIUS (PIPERITES) CIRCELLATUS. *Fries.*
amongst grass. Highgate. Oct. 1863.



milk white, then lilac.


12 M

LACTARIUS (PIPERITES) UVIDUS. *Fries.*
in moist woods. Loughton. Aug. 1872.



milk white, acrid.

LACTARIUS (PIPERITES) FLEXUOSUS. *Fries.*
in pine woods, &c. Hayward Forest, 1881.



milk white, very acrid.




7-10 μ



LACTARIUS (PIPERITES) PYROGALUS. Bull.

in woods, fields, &c. Dinedor, 1881.



milk white, mild.


A



LACTARIUS (PIPERITES) SQUALIDUS. *Krombh.*
in mossy places. *Glamis.*

B


odour pungent.



milk very acrid, white.

M.J.B.


LACTARIUS (PIPERITES) SCOTICUS. *Berk. & Br.*
amongst moss. *Aboyne.*



milk white, peppery.




LACTARIUS (PIPERITES) CAPSICUM. Schulz.
in grassy places. Dumfries, Sept. 1883.



milk acrid, yellow.




LACTARIUS (PIPERITES) CHRYSORHEUS. *Fries.*
in beech woods. Black Park, Langley, 1881.



*stinking, acrid.
milk white, then reddish.*




LACTARIUS (PIPERITES) ACRIS. Bolton.
in woods. Reigate Hill.



milk white, acrid.



LACTARIUS (PIPERITES) UMBRINUS. *Pers.*
in damp places. *Epping Forest.*




milk white, acrid.



7 X 5-6 M

LACTARIUS (PIPERITES) PARGAMENUS. Swartz.
on the ground. Hayward Forest, Sept. 1882.



milk copious, white, acrid.




LACTARIUS (PIPERITES) PIPERATUS. Scop.
on the ground in woods. Common.

milk sparing, white, acrid.



LACTARIUS (PIPERITES) VELLEREUS. *Fries.*
on ground in woods. Hayward Forest, 1883.



pileus pubescent.


no apparent milk.

8-9 μ

LACTARIUS (PIPERITES) EXSUCCUS. Otto.
on the ground. Epping Forest.



LACTARIUS (*DAPETES*) DELICIOSUS. Linn.
 under firs. Lyndhurst, New Forest, 1882.




milk mild, white.



LACTARIUS (RUSSULARIA) PALLIDUS. Pers.


in beech woods. Dinedor, 1881.



milk white, mild.


10-12 M.

LACTARIUS (RUSSULARIA) QUIETUS. *Fries.*
in woods. Chingford, Essex. Nov. 1882.



milk white, peppery.

LACTARIUS (*RUSSULARIA*) AURANTIACUS. *Fries.*
in woods. Woodman's Glade, Epping. Sept. 1888.




milk watery white, mild.

pileus punctate.

10 M

G.M.

LACTARIUS (RUSSULARIA) CREMOR. *Fries, var. PAUPER. Karst.*
in woods, &c. Carlisle, Sept. 1887.




milk white, then greyish.




LACTARIUS (RUSSULARIA) VIETUS. *Fries.*
on the ground. near Hereford, 1871.

odour of bugs.



milk white.


LACTARIUS (RUSSULARIA) CYATHULA. *Fries.*
under birch. Aug. 1862.




milk white, mild.




LACTARIUS (*RUSSULARIA*) CYATHULA. *Fries.*
under birches. Minstead, New Forest. Aug. 1888.





LACTARIUS (RUSSULARIA) RUFUS. Scop.
in pine woods. near Abergele.




milk watery white, slightly acid.

10-12 *M*

odour faint.

LACTARIUS (RUSSULARIA) HELVUS. *Fries.*

amongst moss in moist woods.



milk white, slightly acrid.

pileus woolly.


8-9 M

LACTARIUS (RUSSULARIA) TOMENTOSUS. Otto.
in damp places. Orton Moss, Carlisle, 1886.



milk white, acrid.

LACTARIUS (RUSSULARIA) MAMMOSUS. var. MONSTROSUS. *Fries Icon.*
on the ground in woods. Scarborough, Oct. 1884.




milk mild, white.



6-10M

fragrant.

LACTARIUS (RUSSULARIA) GLYCIOSMUS. *Fries.*
on the ground. *Dinmore, Sept. 1883.*




milk white, then saffron.

10-12 M

LACTARIUS (RUSSULARIA) FULIGINOSUS. *Fries.*

in shady woods. Epping Forest, 1882.






milk white, acrid.



LACTARIUS (RUSSULARIA) PICINUS. *Fries.*
in swampy ground. Forley, 1879.




milk white, acrid.

A




LACTARIUS (*RUSSULARIA*) LILACINUS. *Lasch.*
in woods. Sunny Gutter, Herefordshire, Oct. 1882.

B



H.G.B.


LACTARIUS (*RUSSULARIA*) SPINOSULUS. *Quel. var. VIOLACEUS. Cke.*
on the ground. Chatsworth, Sept. 1883.



milk white, sweet.



LACTARIUS (*RUSSULARIA*) VOLEMUS. *Fries.*
in woods, &c. Breinton, Aug. 1873.




milk white, sweet.



10 X 8 M

LACTARIUS (RUSSULARIA) ICHORATUS. *Batsch.*
in woods, &c. near Hereford.




milk insipid, watery.




7-8M

LACTARIUS (RUSSULARIA) SERIFLUUS. *De Cand.*
in damp places. Haywood Forest, 1882.



milk white, mild.

LACTARIUS (RUSSULARIA) MITISSIMUS. *Fries.*
in woods, &c. Dinedor Common, Oct. 1875.



milk white, mild.



LACTARIUS (*RUSSULARIA*) SUBDULCIS. Bull.
in woods. Epping Forest.



odour fragrant when dried.
milk mild, white.

A

9M

W.P.


LACTARIUS (RUSSULARIA) CAMPHORATUS. Bull.
in woods. The Wrekin. Sept. 1875.

milk rather acrid, white.

B

odour of bugs.

LACTARIUS (RUSSULARIA) CIMICARIUS. Batsch.
amongst grass. Kew Gardens, 1882.



milk watery, whitish, mild.

A




G.M.

LACTARIUS (RUSSULARIA) SUBUMBONATUS. *Lind.*


in woods. Chingford, Epping Forest, 1886.

B



LACTARIUS (RUSSULARIA) MINIMUS. *Smith.*

in fields, &c. Abergavenny.



milk white, mild.




M.J.B.

LACTARIUS (RUSSULARIA) OBNUBILIS. *Lasch.*
in woods.



milk white.


LACTARIUS (PLEUROPUS) OBLIQUUS. *Fries.*
on banks. Dinmore.







RUSSULA (COMPACTÆ) NIGRICANS. Bull.
on the ground. Arboretum, Kew. July, 1888.






RUSSULA (COMPACTÆ) ALBO-NIGRA. *Kromb.*
amongst grass. *Arboretum, Kew.* Aug 1886.






RUSSULA (COMPACTÆ) ADUSTA. Fries.
 under trees. Minstead, New Forest. Aug. 1888.






RUSSULA (COMPACTÆ) DENSIFOLIA. Gillet.
under trees, amongst grass. Kew, Aug. 1886.





mild.

RUSSULA (COMPACTÆ) SEMICREMA. *Fries.*
on the ground. Neasden. Sept. 1881.




*smooth, shining.*8-10 X 6-8 μ .RUSSULA (COMPACTÆ) DELICA. *Fries.**on the ground. Kenilworth, and Neasden. Oct. 1882.*






RUSSULA (COMPACTÆ) MUSTELINA. *Fries.*
in woods. Slough, and Kew Gardens.




mild.




11-12. u.

RUSSULA (*FURCATÆ*) OLIVASCENS. *Fries.*
on the ground. Pleasure Grounds, Kew. Aug. 1886.




at first mild.



*gills crowded,
furcate.*


9.4

RUSSULA (FURCATÆ) FURCATA. Fries.
under trees. Fairmead, Epping, Aug. 1888.




mild.

RUSSULA (FURCATÆ) FURCATÆ. *Fr. var. PICTIPES.*
under trees. Cuckoo Pits, Epping. Aug. 1888.




mild.




9 X 7 M


RUSSULA (*FURCATÆ*) *FURCATÆ*. Fries. var. *OCHROVIRIDIS*.
on the ground. Dinedor. Aug 1882.



peppery.




RUSSULA (*FURCATÆ*) SANGUINEA. *Bull.*
on the ground. Haywood Common.




slowly acrid.

RUSSULA (*FURCATÆ*) ROSACEA. *Fries.*
 amongst grass. *Credinhill Camp.* *Sept. 1882.*





RUSSULA (*FURCATÆ*) MACULATA. Quelet.
 under trees. Woodman's Glade, Epping. Sept. 1888.




rather acrid.



8-10 μ .


spores pale ochre.


RUSSULA (*FURCATÆ*) SARDONIA. *Fries. (variety).*
about larch. Black Park, Langley.



mild.

RUSSULA (*FURCATÆ*) DEPALLENS. Pers.
in woods. Dinedor, Sept. 1881.






mild.

12M


RUSSULA (FURCATÆ) CÆRULEA. Pers.
amongst grass. Minstead, New Forest. Aug. 1888.



intensely acrid.

spores pale ochre.


RUSSULA (FURCATÆ) DRIMEIA. Cooke.
under trees. Black Park, Langley, 1881.



mild.




RUSSULA (*RIGIDÆ*) LACTEA. *Pers.*
on the ground. Credinhill Camp. Sept. 1882.




mealy, mild.

RUSSULA (RIGIDÆ) LACTEA. Pers. var. INCARNATA. Quel.
 under pines, &c. Epping Forest.




mild.



7-8 μ .

RUSSULA (*RIGIDÆ*) VIRESCENS. Schæff.
under trees, Chingford, Epping Forest, Aug. 1884.






RUSSULA (RIGIDÆ) CUTEFRACTA. Cooke.
on the ground. Arboretum, Kew, 1881.



mild.




RUSSULA (RIGIDÆ) CUTEFRACTA. Cooke,
on the ground. Monk's Wood Epping 1881



mild.



RUSSULA (*RIGIDÆ*) LEPIDA. Fries.
under trees. near Morpeth. Aug. 1888.




mild.

RUSSULA (RIGIDÆ) LEPIDA. *Fries.* VARIETY.
amongst grass. *Fairmead, Epping.* Aug. 1888.



acid.

RUSSULA (RIGIDÆ) RUBRA. Fries.
under trees. Monkswood, Epping.




mild.



R. atropurpureus. Krom.

RUSSULA (RIGIDÆ) RUBRA. *Fries. var. SAPIDA.*
under trees. Arboretum, Kew. Sept. 1888.




taste mild, sweet.



RUSSULA (RIGIDÆ) LINNÆI. *Fries.*


on the ground. Crackley Wood, Kenilworth, 1885.




mild.




RUSSULA (RIGIDÆ) XERAMPELINA. Schæff.
amongst grass. Arboretum, Kew.



mild.

9-10 X 7-8 μ

RUSSULA (RIGIDÆ) XERAMPELINA. Schoff,
under trees. Arboretum, Kew. Sept. 1888.




mild.



RUSSULA (RIGIDÆ) OLIVACEA. Schæff.


amongst grass. Pleasure Grounds, Kew. Aug. 1886.



mild.



RUSSULA (HETEROPHYLLÆ) VESCA. Fries.
amongst grass. Arboretum, Kew. 1886.



odour of crab.

A



W.P.

RUSSULA (*RIGIDÆ*) DU PORTII. Phil.
under trees. Mattishall. Sept. 1883.


acrid.



8-9 μ

pileus pruinose.


RUSSULA (*RIGIDÆ*) SEROTINA. Quelet.
under beech trees. Arboretum, Kew. Sept. 1886.




mild.




RUSSULA (HETEROPHYLLÆ) L!LACEA *Quelet.*
under trees. Loughton, Essex. Aug. 1881.




mild.



RUSSULA (HETEROPHYLLÆ) AZUREA. Bres.
amongst grass. Arboretum, Kew, 1888.




mild.



RUSSULA (HETEROPHYLLÆ) CYANOXANTHA.
amongst grass.

Arboretum, Kew.

Schæff. (variety).
Aug. 1888.



mild.



9 X 8 M


RUSSULA (HETEROPHYLLÆ) CYANOXANTHA. Schæff.
under trees. Arboretum, Kew. Aug. 1888.



mild.



RUSSULA (HETEROPHYLLÆ) CYANOXANTHA. Schaff. VARIETY.
 under trees. Arboretum, Kew. Sept. 1888.




mild.

gills crowded, thin.

6-7 M

RUSSULA (HETEROPHYLLÆ) HETEROPHYLLA. *Fries.*
amongst grass. *Arboretum, Kew.* Aug. 1888.




mild.

7-8 M.


gills crowded, narrow.

RUSSULA (HETEROPHYLLÆ) HETEROPHYLLA. *Fries.*
under beech. Arboretum, Kew. Aug. 1886.



mild.

RUSSULA (HETEROPHYLLÆ) GALOCHROA. Bull.
 under trees, Arboretum, Kew Aug. 1888.




peppery.




10x8 M


RUSSULA (HETEROPHYLLÆ) CONSOBRINA. *Fries*
under beech. *Kew Gardens.* Aug. 1886.



very acrid.




RUSSULA (HETEROPHYLLÆ) CONSOBRINA. *Fr. var. INTERMEDIA*
under trees. Arboretum, Kew. July, 1888.



acrid.

RUSSULA (HETEROPHYLLÆ) CONSOBRINA. *Fries. var. SORORIA.*
amongst grass. Kew Gardens. July, 1884.




fœtid, acrid.



8-9 M.

RUSSULA (HETEROPHYLLÆ) FÆTENS. Pers.

under trees. Black Park, Langley. Sept. 1881.




*somewhat acrid,
odour unpleasant.*




10-8 M

whole plant tough and elastic.


RUSSULA (HETEROPHYLLÆ) SUBCÆTENS. *Smith.*
amongst grass. *Pleasure Grounds, Kew. Aug. 1886.*




acid.




RUSSULA (HETEROPHYLLÆ) FELLEA. Fries.
under trees. Arboretum, Kew. Sept. 1881.



becoming acrid.




RUSSULA (HETEROPHYLLÆ) ELEGANS. *Bresadola.*
amongst grass. *Arboretum, Kew.* *Sept. 1886.*



acid.

stem mealy.

RUSSULA (HETEROPHYLLÆ) QUELETHII. *Fries.*
under firs. Black Park, Langley.



acrid.



RUSSULA (HETEROPHYLLÆ) EXPALLENS. Gillet.
under trees. Arboretum, Kew, Oct. 1886.




acrid.



sp. 8μ

RUSSULA (*FRAGILES*) EMETICA. *Fries.*
under trees in woods. *Neasden, Sept. 1881.*




acid.



RUSSULA (FRAGILES) CLUSII. Fries.


under trees. (A) Epping Forest. (B) Black Park, 1881.



fragile, acrid.




RUSSULA (*FRAGILES*) FALLAX Schæff.
on the ground. Neasden. Sept. 1881.




slightly acrid.

RUSSULA (*FRAGILES*) PECTINATA. *Bull.*
on the ground. Breinton, Hereford, 1881.




acrid.

RUSSULA (FRAGILES) OCHROLEUCA. Pers.
on the ground. Arboretum, Kew, 1881.




acid.



disc and stem granulate.


RUSSULA (FURCATÆ) GRANULOSA. Coul.
under trees, Arboretum, Kew. Aug. 1888



mild.




RUSSULA (FRAGILES) ÆRUGINEA. Fries.
on the ground. Kray: Langley.



acrid.




RUSSULA (FRAGILES) FRAGILIS *Fries.*
on the ground. *Epping Forest.*




acrid.

A




RUSSULA (*FRAGILES*) FRAGILIS. *Fries. var. VIOLACEA. Q.*
in woods. *Monks Wood, Epping, 1887.*


B



RUSSULA (*FRAGILES*) FRAGILIS. *Fr. var. NIVEUS. Pers.*
under trees. *Kew Gardens, 1886.*




mild.



9-7 M.

RUSSULA (FRAGILES) FINGIBILIS. Britz.
under trees. Arboretum, Kew. July, 1882.




mild, then slightly acrid.




gills furcate at base.

RUSSULA (FRAGILES) CITRINA. Gillet.
on ground in woods. Black Park. Sept. 1881.



mild.

RUSSULA (FRAGILES) PUNCTATA. *Gillet, var. LEUCOPUS.*
amongst grass. *Arboretum, Kew. July, 1888.*




acrid.




RUSSULA (FRAGILES) VETERIOSA. Fries.

amongst grass under trees. Pleasure Grounds, Kew, 1886.



acrid.


RUSSULA (FRAGILES) VETERINOSA. Fries.
under trees. Arboretum, Kew. Sept. 1888.



mild.




RUSSULA (*FRAGILES*) ROSEIPES. *Secr.* VARIETY.
amongst grass. *Morpeth.* Sept. 1888.




viscid, mild.


RUSSULA (FRAGILES) INTEGRA. *Linn.*
under trees. Arboretum, Kew.



mild.




RUSSULA (FRAGILES) INTEGRA. Linn.
on the ground. Fairmead, Epping, 1882.



mild.



RUSSULA (*FRAGILES*) INTEGRAL. *Linn. var. ALBA.*
under trees. Sutton Park, Warwickshire.




mild.



RUSSULA (*FRAGILES*) DECOLORANS. *Fries.*

in pine woods. Aboyne, 1872.




mild.




12 X 10M

RUSSULA (FRAGILES) BARLÆ. *Quelet.*
amongst grass. *Arboretum, Kew. July, 1888.*






RUSSULA (FRAGILES) AURATA. *With.*
in woods. Horsham.




1097.




12 X 10M.

RUSSULA (FRAGILES) NITIDA. *Fries. var. B Secr. 496.*
in woods. Carlisle. Aug. 1888.




strong scented.



RUSSULA (*FRAGILES*) NITIDA. Pers.

in woods. (A) *Laxton, Norths.* (B) *Arboreium, Kew.*






A



C.B.

*RUSSULA (FRAGILES) PULCHRALIS. Britz.
on the ground. Stapleton Park, 1884.*



B




8-10 μ




mild.

G.M.


*RUSSULA (FRAGILES) NITIDA. Fr. var. CUPRÆA. Krom.
under firs. Bramham Park, Leeds, 1888.*



very fragile.

RUSSULA (*FRAGILES*) ARMENIACA. Cooke.

amongst grass, under trees. Fairmead, Epping. Aug. 1888.




mild.



9 X 8 M

turning foxy where touched.


RUSSULA (*FRAGILES*) PUELLARIS. *Fries.*
under birch. *Morpeth, Sept. 1888.*






turning foxy where touched.

RUSSULA (FRAGILES) PUELLARIS. Fries. VARIETY.
under trees in woods. Morpeth, Sept. 1888.






RUSSULA (FRAGILES) ALUTACEA. *Fries.*
amongst trees. *Minstead, New Forest. Aug. 1888.*



mild



RUSSULA (FRAGILES) ALUTACEA. *Fries.*
on the ground. Epping Forest, 1881.




mild.



12 μ .

gills furcate at the base.


RUSSULA (FRAGILES) OCHRACEA. *Alb. & Schw.*
amongst grass. *Pleasure Grounds, Kew.* Aug. 1886.



mild.




RUSSULA (FRAGILES) LUTEA. Hudson.
in woody places. Hayward Forest, 1881.





RUSSULA (*FRAGILES*) NAUSEOSA. *Fries.*
on the ground. Morpeth, 1889.



strong scented.

A




M.J.B.


RUSSULA (FRAGILES) NAUSEOSA. Fr. var. FLAVIDA.
on the ground. Apethorpe.

strong scented, mild.

B



RUSSULA (FRAGILES) VITELLINA. Fries.
on the ground. Chingford, Essex. Aug. 1886.





mild.

G.M.




RUSSULA (FRAGILES) CHAMELEONTINA. Fries.
 under beeches. Arboretum, Kew, 1886.





CANTHARELLUS CIBARIUS. Fries.
on the ground. Epping Forest, 1886.







CANTHARELLUS CIBARIUS. *Fr. var. RUFIPES. Gill.*
on the ground. Reading, Oct. 1872.




CANTHARELLUS FRIESII *Quelet*
in woods. Sydenham.





CANTHARELLUS AURANTIACUS Fries.
amongst grass. Forest of Dean, 1887.



A





CANTHARELLUS BROWNII. *B. & Br.*
in grassy places. *Hitchin.*

B




CANTHARELLUS UMBONATUS. *Fries.*
amongst moss. *Mossburnford.*






CANTHARELLUS CARBONARIUS *Fries.*
on charred ground. Arborelum. Kew. Nov. 1880




A




CANTHARELLUS ALBIDUS. *Fries.*
on the ground. Fairmead, Epping.

B




CANTHARELLUS HOUGHTONI. *Phillips.*
on the ground, near Hereford.






CANTHARELLUS TUBÆFORMIS. *Fries.*
on the ground. King's Cliffe.






CANTHARELLUS INFUNDIBULIFORMIS. *Fries.*
on the ground. Carlisle, Oct. 1887.





CANTHARELLUS CINEREUS. *Fries.*
on the ground. *Coed Coch.*



CANTHARELLUS CUPULATUS. *Fries.*
on the ground. *near Bristol, Nov. 1882.*






CANTHARELLUS LEUCOPHÆUS. *Novel.*
on the ground. King's Cliffe.



CANTHARELLUS STEVENSONI. *B. & Br.*
on stumps. Glamis, N.B.



A




CANTHARELLUS MUSCIGENUS. *Bull.*
on mosses. *Aboyne.*


B



CANTHARELLUS GLAUCUS. *Batsch.*
on sandy slopes. *near Shrewsbury.*




A

10 X 8-9 μ


M.J.B.


CANTHARELLUS RETIRUGUS. *Fries.*
on mosses. King's Cliffe.

B



CANTHARELLUS LOBATUS. *Fries.*
in swamps. Apethorpe







NYCTALIS CALIGINOSA. *Smith.*
amongst leaves. *Bishop's Wood, Highgate.*





NYCTALIS ASTEROPHORA. *Fries.*
on dead *Russula*. *Epping Forest.*







NYCTALIS PARASITICA. Fries.
on decaying fungi. Hainault Forest.






MARASMIUS (*SCORTEI*) URENS. *Bull*
amongst grass. Pleasure Grounds, Kew. Sept. 1881.






MARASMIUS (*SCORTEI*) PERONATUS. Fries.
 on the ground. Hainault Forest.



odour of garlic.




MARASMIUS (SCORTEI) PORREUS. Pers.
amongst dead leaves. Collyweston.






8 X 5 M

MARASMIUS (COLLYBIA) OREADES. *Fries.*
on lawns, &c. *Kew Gardens.*



A




MARASMIUS (SCORTEI) PLANCUS. *Fries.*
on the ground. Coed Coch, 1882.

B



8 X 6 M


MARASMIUS (SCORTEI) SCORTEUS. *Fries.*
in moist woods. Laxton.



odour of garlic.

G.M.

MARASMIUS (TERGINI) PRASIOSMUS. *Fries.*
amongst leaves. Scarborough.




A



MARASMIUS (TERGINI) VARICOSUS. *Fries.*
in moist places, amongst moss.

B




W.J.B.

MARASMIUS (TERGINI) FUSCO-PURPUREUS. *Pers.*
amongst twigs and leaves. *Apethorpe.*







MARASMIUS (TERGINI) TERGINUS. *Fries.*
amongst leaves. near Bristol.



MARASMIUS (TERGINI) ARCHYROPUS. *Fries.*
amongst leaves. near Bristol.




A




MARASMIUS (TERGINI) WYNNEI. B. & Br.
on the ground. Coed Coch.

B



MARASMIUS (TERGINI) ERYTHROPUS. Pers.
amongst leaves. Fineshade, Nolts.




*stem velvety.***A**

MARASMIUS (TERGINI) TORQUESCENS. *Quelet.*
amongst twigs in moist woods.

*foetid.**stem velvety.***B****8 X 4-5 μ**

MARASMIUS (TERGINI) IMPUDICUS. *Fries.*
about fir trees. *Hanham, Wilts.*




strong scented.

A




MARASMIUS (CALOPODES) SCORODONIUS. *Fries.*
on twigs, &c. near Hereford.

B




MARASMIUS (CALOPODES) CALOPUS. *Pers.*
on twigs, &c. near Bristol.






MARASMIUS (CALOPODES) VAILLANTII. *Fries.*
on twigs and leaves.



MARASMIUS (CALOPODES) ANGULATUS. *Pers.*
on submerged rushes. *Exter, Aug 1884.*



MARASMIUS (CALOPODES) LANGUIDUS. *Lasch.*
on twigs, &c. *Frome.*




A

G.M.

MARASMIUS (CALOPODES) FETIDUS. Sow.
on branches. Pontrilas, 1885.

B

MARASMIUS (CHORDALES) CAUTICINALIS. Wtch.
amongst pine leaves, &c. near Hereford



A

C

4 X 2 M

4 X 3 M


MARASMIUS AMADELPHUS. *Bull.*
on twigs. *Bath.*

MARASMIUS CANDIDUS. *Fries.*
after *Bolton.*

B

4 X 2 M

MARASMIUS (CALOPODES) RAMEALIS. *Bull.*
on bramble, &c. *Chingford, Essex.* *Oct. 1882.*



odour of garlic.

A

16-18 X 10 M

MARASMIUS (MYCENA)


ALLIACEUS. Jacq.
amongst leaves.


B

12 X 6 M


H.G.B.

MARASMIUS (MYCENA) COHÆRENS. Fries.
on the ground. Downton.







MARASMIUS (ROTULÆ) ROTULA. Scop.
on dead twigs. King's Cliffe.




MARASMIUS (ROTULÆ) GRAMINUM. Lib.
on grass, &c. Fineshade, Norths.



MARASMIUS (ROTULÆ) ANDROSACEUS. Bull.
amongst leaves. Epping Forest.






MARASMIUS (ROTULÆ) SPLACHNOIDES. *Fries.*
among pine leaves. *Cabalva.*



MARASMIUS (ROTULÆ) CURREYI. *B. & Br.*
on grass. *Fineshade, Notts.*



MARASMIUS (ROTULÆ) PERFORANS. *Fries.*
on larch leaves. *Coed Coch.*

✓

~~✗~~ ✗ ✗

0


11

11

✗

✗

M.J.B.




MARASMIUS (ROTULÆ) INSITITIUS. *Fries.*
amongst leaves, &c. *King's Cliffe.*

B


5 x 3 M



MARASMIUS (ROTULÆ) HUDSONI. *Pers*
on dead holly leaves *Higghate.*




H.G.B.




MARASMIUS (ROTULÆ) EPICHLŒ. *Fries.*
on dead grass. near Hereford.

G.M.





MARASMIUS (ROTULÆ) ACTINOPHORUS. *B. & Br.*
on naked soil. Kew Gardens, Aug. 1886.

C




MARASMIUS (ROTULÆ) SACCHARINUS. *Batsch.*
on dead leaves, &c. Kenilworth.






MARASMIUS (*ROTULÆ*) EPIPHYLLUS. *Fries.*
on dead leaves. *Epping.*




MARASMIUS (*ROTULÆ*) POLYADELPHUS. *Lasch.*
on dead leaves, &c. *Dinmore.*




= *M. Broomei.* *B.*


W.G.S.

MARASMIUS (*APUS*) SPODOLEUCUS. *B. & Br.*
on elm twigs. *Batheaston.*






LENTINUS TIGRINUS. *Fries.*
on fir in greenhouse. King's Lynn. Aug. 1884.



A



6-7 x 3m

H.G.B.

LENTINUS TIGRINUS. *Fries.*
on wood. *Reigate, July, 1875.*

B



M.J.B.

LENTINUS DUNALII. *Fries.*
on ash. *Dorsetshire.*







7 x 3M

LENTINUS LEPIDEUS. *Fries.*


on dead wood. *Mattishall, Norfolk. May, 1881.*






P. B. Mason


LENTINUS LEPIDEUS. *Fries. (monstrous form.)*
on wood. Brewery. Burton. Mar. 1883.






odour of anise.

LENTINUS COCHLEATUS, Fries.
on stumps. Rotherwas.




LENTINUS VULPINUS, Fries.
on stumps near Hereford.





LENTINUS SCOTICUS. B. & Br.
on wood. Glamis, Nov. 1881.



A



Currey.

LENTINUS FIMBRIATUS. *Currey.*
on log in water.

B



Bolton.

LENTINUS FLABELLIFORMIS. *Bolton.*
on old tree. *Mixenden Mill, 1790.*




PANUS CONCHATUS. *Fries.*
on stumps.



H.G.B.

PANUS TORULOSUS. *Fries.*
on rotten wood. near Hereford.






PANUS STYPTICUS. *Fries.*
on stumps. Downton, 1889.



PANUS FARINACEUS. *Fries.*
on branches. Epping Forest.



PANUS PATELLARIS. *Fries.*
on cherry. Forres, N.B.



A



CANTHARELLUS DEVEXUS. *Fries.*
amongst leaves. *Lyndhurst Road.* *Oct. 1889.*

B




Sowerby.

XEROTUS DEGENER. *Fries.*
on the naked ground.



A



4-5 M


TROGIA CRISPA. Fries.
on branches. Morpeth. Dec. 1883.

B



6 X 4 M

SCHIZOPHYLLUM COMMUNE. Fries.
on logs. near Ely.






A

4 X 2 μ


LENZITES BETULINA. *Fries.*
on stumps. *Epping Forest.*



B

7 X 5 μ

LENZITES FLACCIDA. *Bull.*
on beech. *Hainault Forest.*





A

LENZITES SEPIARIA. *Fries.*
on fir trunks. Northamptonshire.



B


7-8 x 4m

LENZITES ABIETINA. *Fries.*
on fir trunks and branches. Glasgow.








RUSSULA (FRAGILES) NAUSEOSA. Fries.
on the ground. Morpeth, 1889.











QK Cooke, Mordecai Cubitt
607 Illustrations of British
C67 Fungi
v.7

Biomed

**PLEASE DO NOT REMOVE
CARDS OR SLIPS FROM THIS POCKET**

UNIVERSITY OF TORONTO LIBRARY
